

Annals of the Rheumatic Diseases

Leader

- 241 <http://www.ilar.org> A source of information for rheumatologists, allied health professionals, medical students and the general public *R Armstrong, J J (Hans) Rasker*

Review

- 243 Cytokines and systemic lupus erythematosus *G S Dean, J Tyrrell-Price, E Crawley, D A Isenberg*

Case studies in diagnostic imaging

- 252 A self limiting tumour *S T Quek, A Unger, V N Cassar-Pullicino, S N J Roberts*

Extended reports

- 257 Sequential changes of KL-6 in sera of patients with interstitial pneumonia associated with polymyositis/dermatomyositis *S Bandoh, J Fujita, Y Ueda, S Hojo, M Tokuda, H Dobashi, N Kurata, T Yoshinouchi, Y Ohtsuki, N Kohno, J Takahara*
- 263 Differential Th1/Th2 cytokine patterns in chronic arthritis: interferon γ is highly expressed in synovium of rheumatoid arthritis compared with seronegative spondyloarthropathies *J D Cañete, S E Martínez, J Farrés, R Sanmartí, M Blay, A Gómez, G Salvador, J Muñoz-Gómez*
- 269 Vertebral fractures and bone mineral density in idiopathic, secondary and corticosteroid associated osteoporosis in men *S F Evans, M W J Davie*
- 276 Psychological, behavioural, and social adjustment in children and adolescents with juvenile chronic arthritis *A C J Huygen, W Kuis, G Sinnema*
- 283 Monocytes from systemic lupus erythematosus patients are severely altered in phenotype and lineage flexibility *F Steinbach, F Henke, B Krause, B Thiele, G-R Burmester, F Hiepe*
- 289 Assessing the prevalence of hand osteoarthritis in epidemiological studies. The reliability of a radiological hand scale *S Kessler, P Dieppe, J Fuchs, T Stürmer, K P Günther*

Concise reports

- 293 Association of the TNFa13 microsatellite with systemic sclerosis in Japanese patients *F Takeuchi, H Nabeta, M Füssel, K Conrad, K-H Frank*
- 297 Circulating concentrations of soluble L-selectin (CD62L) in patients with primary Sjögren's syndrome *M García-Carrasco, P Pizcueta, R Cervera, M Ramos-Casals, A Sisó, G de la Red, M Ingelmo, J Font, P Engel*
- 300 Prevalence of Chlamydia trachomatis in urine of male patients with ankylosing spondylitis is not increased *M van der Paardt, J C van Denderen, A J C van den Brule, S A Morré, I E van der Horst-Bruinsma, P D Bezemer, B A C Dijkmans*
- 303 Activation of the transcription factor NF- κ B in the rat air pouch model of inflammation *L Ellis, V Gilston, C C-Y Soo, C J Morris, B L Kidd, P G Winyard*
- 308 Bone mineral density and biochemical parameters of bone metabolism in female patients with systemic lupus erythematosus *K Redlich, S Ziegler, H P Kiener, S Spitzauer, P Stohlawetz, P Bernecker, F Kainberger, S Grampp, S Kudlacek, W Woloszczuk, J S Smolen, P Pietschmann*
- 311 Response to methotrexate in early rheumatoid arthritis is associated with a decrease of T cell derived tumour necrosis factor α , increase of interleukin 10, and predicted by the initial concentration of interleukin 4 *M Rudwaleit, Z Yin, S Siegert, M Grolms, A Radbruch, J Braun, J Sieper*
- 315 Urokinase-mediated fibrinolysis in the synovial fluid of rheumatoid arthritis patients may be affected by the inactivation of single chain urokinase type plasminogen activator by thrombin *E A M Braat, A F H Jie, H K Rondan, B Beekman, D C Rijken*

Correspondence

- 319 Cutaneous leucocytoclastic vasculitis caused by cyclosporin A (Sandimmun) *M N Gupta, R D Sturrock, G Gupta*
- 319 An evidence based EULAR meeting? *P Dieppe*
- 319 Psychosocial risks for low back pain: are these related to work? *D A Fishbain*

This issue live and online
www.annrheumdis.com