Could we use salivary gland ultrasonography as a prognostic marker in Sjogren's syndrome? Response to: 'Ultrasonographic damages of major salivary glands are associated with cryoglobulinemic vasculitis and lymphoma in primary Sjogren's syndrome: are the ultrasonographic features of the salivary glands new prognostic markers in Sjogren's syndrome?' by Coiffier *et al*

We thank Dr Coiffier and colleagues for their interesting letter in which they suggest that damages of major salivary glands (SG) might be associated with cryoglobulinemic vasculitis and lymphoma in primary Sjögren's syndrome. The authors raise the question whether the ultrasonographic (US) features of the salivary glands [our proposed novel ultrasound scoring system] could be used as the new prognostic markers for cryoglobulinemic vasculitis and lymphoma in patients with Sjögren's syndrome.¹

In their recent study of different US scoring systems in 97 sicca syndrome patients, Coiffier et al found three B-cell lymphomas in the primary Sjogren's syndrome (pSS) group and three, two, and one patient with cryoglobulinemic vasculitis in the pSS, secondary Sjogren's syndrome (sSS) and control groups, respectively.² The authors concluded that the detection of B-cell lymphomas or cryoglobulinemic vasculitis was associated with pathological US findings regardless of the scoring used. The reported US pathological features of SG were either numerous cystic lesions without healthy parenchyma or fibrous glands scored as a grade 3 according to the new semi quantitative scoring system described by the SG sub group of the OMERACT US working group.¹ Although their findings may suggest specific ultrasound features as a risk factor for developing lymphoma or cryoglobulinemic vasculitis, we think at this point in time it is premature to draw such a conclusion.³ Ultrasound may reveal predisposing factors, but these are not proven to be pathognomonic of lymphoma. Indeed, several predictors of lymphoma in pSS such as epidemiological, clinical (permanent swelling of the SG, palpable purpura, organomegaly), biological (cryoglobulinaemia, or low complement levels) and histopathological findings should also be taken into account.⁴⁻⁹ Large sample and longitudinal studies assessing these clinical and biological predictors of lymphoma with US are currently ongoing and will probably shed more light on this challenging issue. Furthermore, pSS disease activity, for example, assessed by the EULAR SS index, can be used as a clinical predictor of lymphoma development with a dose effect.¹⁰

SG enlargement (eg, clinical aspects: unilateral, fixed and hard parotid glands) is regarded as the most dominant clinical symptom for lymphoma in patients with pSS.⁶ SGUS performed by well-trained ultrasonographers can provide a precise structural assessment of the glands' surface compared with clinical examination.¹¹ In case of SGUS grade 3, that is, complete destruction of the gland, with numerous hypoechoic or hyper-echoic bands, the detection of abnormal lymph nodes should raise awareness of possible lymphoma development. Suspicion of abnormal lymph nodes can be confirmed during a long term monitoring of pSS patients especially those with high risk of lymphoma development. In addition, Doppler assessment of gland's vascularisation in pSS might be of help to detect at risk ultrasound lesions. To this end, the forthcoming results of a longitudinal study for development of consensual Doppler US scoring of gland's vascularisation in pSS can be quite helpful.

Sandrine Jousse-Joulin [©], ¹ Maria Antonietta D'Agostino,² Alojzija Hočevar [©], ³ Esperanza Naredo,⁴ Lene Terslev, ⁵ Sarah Ohrndorf, ⁶ Annamaria Iagnocco, ⁷ Wolfgang A Schmidt, ⁸ Stephanie Finzel, ⁹ Zarrin Alavi, ¹⁰ George A W Bruyn, ^{11,12} on behalf the Salivary Gland subgroup of the OMERACT US working group

¹Rheumatology, Brest University Hospital, Brest, France
²Rheumatology Department, INSERM U1173, Laboratoire d'Excellenc, Paris, France
³Rheumatology, Universitiy Medical Centre Ljubljana, Ljubljana, Slovenia
⁴Rheumatology, Severo Ochoa Hospital, Madrid, Spain

⁵Rheumatology, Glostrup University Hospital, Copenhagen, Denmark ⁶Rheumatology and Clinical Immunology, Charité-Universitätsmedizin Berlin, Berlin, Germany

⁷Scienze Cliniche e Biologiche, Università degli Studi di Torino, Rome, Italy ⁸Rheumatology, Medical Centre for Rheumatology Berlin Buch, Berlin, Germany ⁹Rheumatology and Clinical Immunology, University Medical Center Freiburg, Freiburg, Germany

¹⁰INSERM, CIC 1412, Brest University Hospital, Brest, France

¹¹MC Group Hospitals, Lelystad, The Netherlands

¹²Reumakliniek Flevoland, Lelystad, The Netherlands

Correspondence to Dr Sandrine Jousse-Joulin, Rheumatology Department, Cavale Blanche Hospital and Brest Occidentale University, Brest EA 2216, ERI 29, France; sandrine.jousse-joulin@chu-brest.fr

Handling editor Josef S Smolen

Contributors All co-authors participated in the writing and reading of this correspondence letter.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Jousse-Joulin S, D'Agostino MA, Hočevar A, et al. Ann Rheum Dis 2021;80:e112.

Received 16 September 2019 Accepted 18 September 2019 Published Online First 10 October 2019



http://dx.doi.org/10.1136/annrheumdis-2019-216122

Ann Rheum Dis 2021;80:e112. doi:10.1136/annrheumdis-2019-216327

ORCID iDs

Sandrine Jousse-Joulin http://orcid.org/0000-0002-5479-5887 Alojzija Hočevar http://orcid.org/0000-0002-7361-6549

REFERENCES

- 1 Jousse-Joulin S, D'Agostino MA, Nicolas C, et al. Video clip assessment of a salivary gland ultrasound scoring system in Sjögren's syndrome using consensual definitions: an OMERACT ultrasound Working group reliability exercise. Ann Rheum Dis 2019;78:967–73.
- 2 Coiffier G, Martel A, Albert J-D. Ultrasonographic damages of major salivary glands are associated with cryoglobulinemic vasculitis and lymphoma in primary Sjogren's syndrome: are the ultrasonographic features of the salivary glands new prognostic markers in Sjogren's syndrome? *Ann Rheum Dis* 2021;80:e111.
- 3 Theander E, Vasaitis L, Baecklund E, et al. Lymphoid organisation in labial salivary gland biopsies is a possible predictor for the development of malignant lymphoma in primary Sjogren's syndrome. Ann Rheum Dis 2011;70:1363–8.



Ann Rheum Dis: first published as 10.1136/annrheumdis-2019-216327 on 10 October 2019. Downloaded from http://ard.bmj.com/ on June 10, Erasmushogeschool Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies. 2025 at Department GEZ-LTA

Correspondence response

- 4 Anaya J, McGuff H, Banks P, et al. Clinicopathological factors relating malignant lymphomawith Sjögren's syndrome. *Semin Arthritis Rheum* 1996;25:337–46.
- 5 Quartuccio L, Isola M, Baldini C, et al. Biomarkers of lymphoma in Sjögren's syndrome and evaluation of the lymphoma risk in prelymphomatous conditions: Results of a multicenter study. J Autoimmun 2014;51:75–80.
- 6 Nocturne G, Pontarini E, Bombardieri M, et al. Lymphomas complicating primary Sjögren's syndrome: from autoimmunity to lymphoma. *Rheumatology* 2019;8.
- 7 Nishishinya MB, Pereda CA, Muñoz-Fernández S, et al. Identification of lymphoma predictors in patients with primary Sjögren's syndrome: a systematic literature review and meta-analysis. *Rheumatol Int* 2015;35:17–26.
- 8 Solans-Laqué R, López-Hernandez A, Angel Bosch-Gil J, et al. Risk, predictors, and clinical characteristics of lymphoma development in primary Sjögren's syndrome. Semin Arthritis Rheum 2011;41:415–23.
- 9 Nocturne G, Virone A, Ng W-F, et al. Rheumatoid factor and disease activity are independent predictors of lymphoma in primary Sjögren's syndrome. Arthritis & Rheumatology 2016;68:977–85.
- 10 Nocturne G, Mariette X. Sjögren syndrome-associated lymphomas: an update on pathogenesis and management. *Br J Haematol* 2015;168:317–27.
- 11 Marteau P, Cornec D, Gouillou M, et al. Assessment of major salivary gland size in primary Sjögren's syndrome: comparison between clinical examination and ultrasonography. *Joint Bone Spine* 2019. doi:10.1016/j.jbspin.2019.01.025. [Epub ahead of print: 11 Feb 2019].